



# Workshop on Methane on Mars

Current observations, interpretation and future plans 25-27 November 2009, ESRI, Frascati, Italy

Methane has been detected in the Martian atmosphere by ground-based telescopes and from orbit. This discovery indicates that the planet is either biologically or geologically active. The goal of the workshop is to review the available measurements, the potential reservoirs and release mechanisms of Methane and its circulation in the atmosphere, and to discuss all possible origins of this constituent.



Methane release Northern summer



Methane concentration (ppb) versus longitude (degrees)

Deadline for abstracts: 1<sup>st</sup> September 2009

<http://www.congrex.nl/09c26/>

Scientific Organizing Committee: Nicolas Abetti (ESA/ESAC, Madrid), Susan Abreu (The University of Michigan), Vincent Chevreton (University of Arkansas), Agustín Chicarro (ESA/ESTEC, Noordwijk), Thomas Encrenaz (Observatoire de Paris-Meudon), Enrico Fieschi (ASI, Roma), Vittorio Formisano (IFSI, INF, Frascati), Franck Lelligne (LATMOS, Paris), Paul Mahaffy (NASA Goddard Space Flight Center), Michael Mumma (NASA Goddard Space Flight Center), Richard Pinn (University of Colorado, Berkeley), Olivier Witasse (ESA/ESTEC, Noordwijk). Local Organizing Committee: Claire Bingham (ESA/ESTEC, Noordwijk), Carman Compagno (ESA/ESRI, Italy), Sylvie Esphahan (ASI, Roma), Olivier Witasse (ESA/ESTEC, Noordwijk).



NOTE ADDED BY JPL WEBMASTER: This document was prepared by the European Space Agency. The content has not been approved or adopted by, NASA, JPL, or the California Institute of Technology. This document is being made available for information purposes only, and any views and opinions expressed herein do not necessarily state or reflect those of NASA, JPL, or the California Institute of Technology.

O. Witasse, A. Chicarro, and J. L. Vago

ESA, ESTEC (NL)


# Scientific Organising Committee

Nicolas Altobelli	(ESA/ESAC, Madrid)
Sushil Atreya	(The University of Michigan)
Vincent Chevrier	(University of Arkansas)
Agustin Chicarro	(ESA/ESTEC, Noordwijk)
Therèse Encrenaz	(Observatoire de Paris-Meudon)
Enrico Flamini	(ASI, Roma)
Vittorio Formisano	(IFSI- INAF, Roma)
Franck Lefèvre	(LATMOS, Paris)
Paul Mahaffy	(NASA Goddard Space Flight Center)
Michael Mumma	(NASA Goddard Space Flight Center)
Buford Price	(University of California, Berkeley)
Olivier Witasse	(ESA/ESTEC, Noordwijk)
Richard W. Zurek	(NASA Jet Propulsion Laboratory)

# Some Numbers

- Almost 100 participants (66% Europe, 20% US, and the rest from other places);
- 3 days;
- 13 hours of talks;
- **4.5 hours of discussion;**
- 1.5 hours for posters;
- 5.5 hours breaks (lunch, coffee);
- 2 social events: welcome drink and dinner.

# Joint ESA-ASI Methane Workshop, 27–27 November 2009, Frascati (I)



Excellent talks and motivated speakers



Lively discussions



Tough session chairs



Intense poster sessions

# Sessions


1. Observations of methane from Mars orbit;
2. Earth-based observations of Martian methane;
3. Martian surface and subsurface data, and laboratory measurements relevant to the study of methane;
4. Origin of methane;
5. Storage, release, and delivery of methane;
6. Atmospheric circulation and chemistry;
7. Microbial life, metabolism in water ice, and biological experiments under Martian conditions;
8. Future measurements of methane, related trace gases, and isotopes.

# Some highlights



# Mars Express Observations


Vertical profile of H<sub>2</sub>O, CO, and CH<sub>4</sub> during Spring



Message: Methane is not found close to the surface.

# Earth-based Observations

Mumma & Villanueva



Maximum abundance  
Observed at  $L_s$  220°  
(mid-spring in South)

Maximum abundance  
Observed at  $L_s$  155°  
(late summer in North)

Message: Methane is found mainly over older terrain.




# Summary of the observations

- Mars Express/PFS :
  - New CH<sub>4</sub> vertical profiles;
  - New maps, characterized by large contrasts on regional scales;
  - Maximum abundances of methane occur near the north pole in mid to late summer;
- Earth-based observations:
  - Confirmation of previous findings;
  - CH<sub>4</sub> is found over old terrains;
- MGS/TES observations of methane were also reported, showing an enhancement in methane over Tharsis and Arabia Terra;
- Further work is required to reconcile the spacecraft and ground-based observations.

# Geological and Geochemical Context


Methane-rich regions seem to have landforms/mineralogy with possible implications for past methane-related geological processes. Correlating methane to regional geology is a potentially powerful tool that we are just beginning to explore.



Nilius Fossae (Ehlmann et al 2009)



Isidis Planitia  
Mounds (HiRISE)



Mud volcanoes  
Terrestrial analogue


# Origin of Martian Methane

- The presence of methane on Mars remains puzzling;
- Origin: biological or geochemical (or both);
- All other potential production mechanisms —exogenic (comets, meteorites), volcanic, and atmospheric— have been ruled out;
- If the production of methane is ongoing, it points to the presence of underground aquifers;
- If the methane was produced when Mars was supposedly warmer and liquid water flowed on the surface, it could have been stored (as clathrates) and is possibly being released from time to time.

# Storage

## Release of methane clathrate particles

- 1- Formation of clathrate in the crust at high (lithostatic) pressure.
- 2- Fracturation of subsurface by tectonics, yielding local macroporosity
- 3- Erosion of (friable) clathrate host framework and uplift of particles.
- 4- Release to the atmosphere of (ultra-stable) clathrate particles.



New hypothesis (E. Chassefière): Methane release could come from clathrates in the atmosphere. This could explain the (not well established correlation with H<sub>2</sub>O).



# Atmospheric Circulation

Lefèvre & Forget

## Conclusions

- The "conventional" atmospheric chemistry does not produce measurable methane variations on Mars, even in the case of a current, episodic, and localized source.
- The condensation/sublimation cycle of  $\text{CO}_2$  should generate large-scale methane variations at high latitudes (*but they differ from what is observed*).
- CSHELL/NIRSPEC: In the most favourable case, an atmospheric  $\text{CH}_4$  lifetime of less than 200 days (seasonal release) or  $\sim 2$  Earth years (single event) is necessary to reproduce the observations.
- PFS: measurements at high latitudes require a lifetime of less than  $\sim 3$  Earth years. Longitudinal variations at high latitudes and seasonal trends at mid-to-low latitudes cannot be reproduced.
- The  $\text{CH}_4$  source: quantitative agreement with the observations requires considerable amounts:

~150 000 tonnes  
~ 80 000 tonnes  
~ 50 000 tonnes

CSHELL/NIRSPEC, seasonal release  
CSHELL/NIRSPEC, single event  
PFS, polar summer

Mid-Atlantic Ridge: 50 000-130 000  $\text{t yr}^{-1}$   
(Keir et al., 2005)

# Atmospheric Circulation

Lefèvre & Forget

## Conclusions

### Solutions ?

- fast atmospheric loss of methane by chlorine:
  - is not supported by observations of HCl
- fast atmospheric loss of methane by electrochemistry:
  - is not supported by current observations of CO, H<sub>2</sub>O<sub>2</sub>, and O<sub>3</sub>
- fast loss of methane in the regolith:
  - must be extraordinarily rapid (< 24 h) to satisfy the observations
  - is not supported by current observations of other minor species (CO, H<sub>2</sub>O<sub>2</sub>, O<sub>3</sub>), or must be highly selective
  - is for the moment not supported by laboratory data (Gough et al., this workshop)

→ Observed variations of methane are unexplained by known atmospheric chemistry and physics



# Microbial Life

A fascinating and very educational series of reports on microbial life, metabolism in water ice, and biological experiments under Martian conditions was presented by many authors.

The presentations considered perchlorate-reducing bacteria, microbes trapped in extreme environments, the mechanism and complexity of terrestrial methanogenesis, the extent to which cryoenvironments could be active microbial ecosystems, terrestrial analogs and methanogen energy sources.

# Future Measurements

- Ongoing Mars Express measurements (until at least end of 2012);
- New observations with CRILES on the Very Large Telescope;
- IRTF, Sofia, Herschel;
- MSL: SAM and TLS experiments (from 2012 onwards);
- ExoMars Trace Gas Orbiter (from 2017 onwards);
- ExoMars Rover: MOMA experiment (from 2019 onwards);
- Discussion raised the novel possibility of a satellite at a Mars Lagrange point to provide global imaging of atmospheric composition with very high spectral resolution.

# Presentations

Available on-line at:

<http://sci.esa.int/marsmethane2009>

Including the book of abstracts and some pictures.



# Proceedings

- A special issue is being prepared in *Planetary and Space Science*;
- 25 papers have been submitted;
- To be published after the summer 2010.

# Workshop on Methane on Mars

Current observations, interpretation and future plans 25-27 November 2009, ESRI, Frascati, Italy

Methane has been detected in the Martian atmosphere by ground-based telescopes and from orbit. This discovery indicates that the planet is either biologically or geologically active. The goal of the workshop is to review the available measurements, the potential reservoirs and release mechanisms of Methane and its circulation in the atmosphere, and to discuss all possible origins of this constituent.



Methane release Northern summer



Methane (Terra) (Rising the northern hemisphere)  
Source: Orosi, Trent, Scheraga, NASA



Methane abundance (ppb) as a function of longitude (Mars Eastern ERT)

Deadline for abstracts: 1<sup>st</sup> September 2009

<http://www.congrex.nl/09c26/>

Scientific Organizing Committee: Nicolas Audebert (ESA/ESTEC, Madrid), Susan Atreya (The University of Michigan), Vincent Chevreton (University of Arkansas), Agustín Chicarro (ESA/ESTEC, Noordwijk), Thomas Encrenaz (Observatoire de Paris-Meudon), Enrico Fieschi (ASI, Roma), Vittorio Formisano (IFSI, INF, Frascati), Franck Lelligne (LATMOS, Paris), Paul Mahaffy (NASA Goddard Space Flight Center), Michael Mumma (NASA Goddard Space Flight Center), Richard Pinn (University of Colorado, Berkeley), Olivier Witasse (ESA/ESTEC, Noordwijk). Local Organizing Committee: Claire Bingham (ESA/ESTEC, Noordwijk), Carman Compagno (ESA/ESTEC, Italy), Sylvie Esphahan (ASI, Roma), Olivier Witasse (ESA/ESTEC, Noordwijk).



NOTE ADDED BY JPL WEBMASTER: This document was prepared by the European Space Agency. The content has not been approved or adopted by, NASA, JPL, or the California Institute of Technology. This document is being made available for information purposes only, and any views and opinions expressed herein do not necessarily state or reflect those of NASA, JPL, or the California Institute of Technology.

O. Witasse, A. Chicarro, and J. L. Vago

ESA, ESTEC (NL)

# Scientific Organising Committee


Nicolas Altobelli	(ESA/ESAC, Madrid)
Sushil Atreya	(The University of Michigan)
Vincent Chevrier	(University of Arkansas)
Agustin Chicarro	(ESA/ESTEC, Noordwijk)
Therèse Encrenaz	(Observatoire de Paris-Meudon)
Enrico Flamini	(ASI, Roma)
Vittorio Formisano	(IFSI- INAF, Roma)
Franck Lefèvre	(LATMOS, Paris)
Paul Mahaffy	(NASA Goddard Space Flight Center)
Michael Mumma	(NASA Goddard Space Flight Center)
Buford Price	(University of California, Berkeley)
Olivier Witasse	(ESA/ESTEC, Noordwijk)
Richard W. Zurek	(NASA Jet Propulsion Laboratory)



# Some Numbers

- Almost 100 participants (66% Europe, 20% US, and the rest from other places);
- 3 days;
- 13 hours of talks;
- **4.5 hours of discussion;**
- 1.5 hours for posters;
- 5.5 hours breaks (lunch, coffee);
- 2 social events: welcome drink and dinner.

# Joint ESA-ASI Methane Workshop, 27–27 November 2009, Frascati (I)



Excellent talks and motivated speakers



Lively discussions

Tough session chairs



Intense poster sessions




# Sessions

1. Observations of methane from Mars orbit;
2. Earth-based observations of Martian methane;
3. Martian surface and subsurface data, and laboratory measurements relevant to the study of methane;
4. Origin of methane;
5. Storage, release, and delivery of methane;
6. Atmospheric circulation and chemistry;
7. Microbial life, metabolism in water ice, and biological experiments under Martian conditions;
8. Future measurements of methane, related trace gases, and isotopes.

# Some highlights

# Mars Express Observations


Vertical profile of H<sub>2</sub>O, CO, and CH<sub>4</sub> during Spring



Message: Methane is not found close to the surface.

# Earth-based Observations

Mumma & Villanueva



Maximum abundance  
Observed at  $L_s$  220°  
(mid-spring in South)

Maximum abundance  
Observed at  $L_s$  155°  
(late summer in North)

Message: Methane is found mainly over older terrain.




# Summary of the observations

- Mars Express/PFS :
  - New CH<sub>4</sub> vertical profiles;
  - New maps, characterized by large contrasts on regional scales;
  - Maximum abundances of methane occur near the north pole in mid to late summer;
- Earth-based observations:
  - Confirmation of previous findings;
  - CH<sub>4</sub> is found over old terrains;
- MGS/TES observations of methane were also reported, showing an enhancement in methane over Tharsis and Arabia Terra;
- Further work is required to reconcile the spacecraft and ground-based observations.

# Geological and Geochemical Context


Methane-rich regions seem to have landforms/mineralogy with possible implications for past methane-related geological processes. Correlating methane to regional geology is a potentially powerful tool that we are just beginning to explore.



Nillie Fossae (Ehlmann et al 2009)



Isidis Planitia  
Mounds (HIRISE)



Mud volcanoes  
Terrestrial analogue


# Origin of Martian Methane

- The presence of methane on Mars remains puzzling;
- Origin: biological or geochemical (or both);
- All other potential production mechanisms —exogenic (comets, meteorites), volcanic, and atmospheric— have been ruled out;
- If the production of methane is ongoing, it points to the presence of underground aquifers;
- If the methane was produced when Mars was supposedly warmer and liquid water flowed on the surface, it could have been stored (as clathrates) and is possibly being released from time to time.

# Storage

## Release of methane clathrate particles

- 1- Formation of clathrate in the crust at high (lithostatic) pressure.
- 2- Fracturation of subsurface by tectonics, yielding local macroporosity
- 3- Erosion of (friable) clathrate host framework and uplift of particles.
- 4- Release to the atmosphere of (ultra-stable) clathrate particles.



New hypothesis (E. Chassefière): Methane release could come from clathrates in the atmosphere. This could explain the (not well established correlation with H<sub>2</sub>O).



# Atmospheric Circulation

Lefèvre & Forget

## Conclusions

- The "conventional" atmospheric chemistry does not produce measurable methane variations on Mars, even in the case of a current, episodic, and localized source.
- The condensation/sublimation cycle of  $\text{CO}_2$  should generate large-scale methane variations at high latitudes (*but they differ from what is observed*).
- CSHELL/NIRSPEC: In the most favourable case, an atmospheric  $\text{CH}_4$  lifetime of less than 200 days (seasonal release) or  $\sim 2$  Earth years (single event) is necessary to reproduce the observations.
- PFS: measurements at high latitudes require a lifetime of less than  $\sim 3$  Earth years. Longitudinal variations at high latitudes and seasonal trends at mid-to-low latitudes cannot be reproduced.
- The  $\text{CH}_4$  source: quantitative agreement with the observations requires considerable amounts:

$\sim 150\,000$  tonnes  
 $\sim 80\,000$  tonnes  
 $\sim 50\,000$  tonnes

CSHELL/NIRSPEC, seasonal release  
CSHELL/NIRSPEC, single event  
PFS, polar summer

Mid-Atlantic Ridge:  $50\,000\text{--}130\,000\text{ t yr}^{-1}$   
(Keir et al., 2005)

# Atmospheric Circulation

Lefèvre & Forget

## Conclusions

### Solutions ?

- fast atmospheric loss of methane by chlorine:
  - is not supported by observations of HCl
- fast atmospheric loss of methane by electrochemistry:
  - is not supported by current observations of CO, H<sub>2</sub>O<sub>2</sub>, and O<sub>3</sub>
- fast loss of methane in the regolith:
  - must be extraordinarily rapid (< 24 h) to satisfy the observations
  - is not supported by current observations of other minor species (CO, H<sub>2</sub>O<sub>2</sub>, O<sub>3</sub>), or must be highly selective
  - is for the moment not supported by laboratory data (Gough et al., this workshop)

→ Observed variations of methane are unexplained by known atmospheric chemistry and physics



# Microbial Life

A fascinating and very educational series of reports on microbial life, metabolism in water ice, and biological experiments under Martian conditions was presented by many authors.

The presentations considered perchlorate-reducing bacteria, microbes trapped in extreme environments, the mechanism and complexity of terrestrial methanogenesis, the extent to which cryoenvironments could be active microbial ecosystems, terrestrial analogs and methanogen energy sources.

# Future Measurements

- Ongoing Mars Express measurements (until at least end of 2012);
- New observations with CRILES on the Very Large Telescope;
- IRTF, Sofia, Herschel;
- MSL: SAM and TLS experiments (from 2012 onwards);
- ExoMars Trace Gas Orbiter (from 2017 onwards);
- ExoMars Rover: MOMA experiment (from 2019 onwards);
- Discussion raised the novel possibility of a satellite at a Mars Lagrange point to provide global imaging of atmospheric composition with very high spectral resolution.

# Presentations

Available on-line at:

<http://sci.esa.int/marsmethane2009>

Including the book of abstracts and some pictures.

# Proceedings

- A special issue is being prepared in *Planetary and Space Science*;
- 25 papers have been submitted;
- To be published after the summer 2010.